


FLUX OF LARGE METEOROIDS FROM LUNAR IMPACT MONITORING AND INFRASOUND

Bill Cooke


Lead, NASA Meteoroid Environments Office

william.j.cooke@nasa.gov

Meteoroid - small rocky/
icy debris out in space




Meteoroid - small rocky/
icy debris out in space



Meteor - streak of light/ionization
produced by a meteoroid ablating in
an atmosphere

Meteoroid - small rocky/
icy debris out in space



Meteor - streak of light/ionization
produced by a meteoroid ablating in
an atmosphere

Fireball - meteor with a peak
brightness greater than the
planet Venus ($m_v -4$). Also
called a bolide





Super bolide - a meteor sufficiently energetic to be detected by seismic/other sensors ($m_v \sim -18$)



Super bolide - a meteor sufficiently energetic to be detected by seismic/other sensors ($m_v \sim -18$)



Meteorite - any part of a meteoroid/asteroid that makes it to the surface

ASTEROID OR METEOROID?

Size is the discriminator, but no “official” boundary

ASTEROID OR METEOROID?

Size is the discriminator, but no “official” boundary

Some scientists say meteoroids are smaller than one meter (beach ball size); others place the limit at 10 meters (garage size)


ASTEROID OR METEOROID?

Size is the discriminator, but no “official” boundary

Some scientists say meteoroids are smaller than one meter (beach ball size); others place the limit at 10 meters (garage size)

International Astronomical Union definition: “A meteoroid is a solid object moving in interplanetary space, of a size considerably smaller than an asteroid and considerably larger than an atom.”

How NASA handles meteoroids, asteroids, and space junk




Earth-approaching asteroids




Near Earth
Object Office
(JPL)

How NASA handles meteoroids, asteroids, and space junk



Earth-approaching asteroids


Near Earth
Object Office
(JPL)



Man-made orbital debris

Orbital Debris
Program
Office (JSC)

How NASA handles meteoroids, asteroids, and space junk




Earth-approaching asteroids

Near Earth
Object Office
(JPL)



Meteoroids and meteors

Meteoroid
Environment
Office (MSFC)



Man-made orbital debris

Orbital Debris
Program
Office (JSC)

What does the Meteoroid Environments Office (MEO) do?

What does the Meteoroid Environments Office (MEO) do?

Develop, maintain, and distribute an sporadic meteoroid model suitable for spacecraft engineering/design

What does the Meteoroid Environments Office (MEO) do?

Develop, maintain, and distribute an sporadic meteoroid model suitable for spacecraft engineering/design

Provide meteor shower forecasts to NASA/USG spacecraft operators

What does the Meteoroid Environments Office (MEO) do?

Develop, maintain, and distribute an sporadic meteoroid model suitable for spacecraft engineering/design

Provide meteor shower forecasts to NASA/USG spacecraft operators

Monitor the meteoroid environment in near-Earth space (SSA)

What does the Meteoroid Environments Office (MEO) do?

Develop, maintain, and distribute an sporadic meteoroid model suitable for spacecraft engineering/design

Provide meteor shower forecasts to NASA/USG spacecraft operators

Monitor the meteoroid environment in near-Earth space (SSA)

Conduct and manage research to improve sporadic and shower meteoroid models

MAY CAMELOPARDALIDS

On May 24, 2014 Earth will encounter multiple streams of debris laid down by Comet 209P LINEAR

MAY CAMELOPARDALIDS

On May 24, 2014 Earth will encounter multiple streams of debris laid down by Comet 209P LINEAR

This will produce a new meteor shower, never before seen

MAY CAMELOPARDALIDS

On May 24, 2014 Earth will encounter multiple streams of debris laid down by Comet 209P LINEAR

This will produce a new meteor shower, never before seen

Rates predicted to be from 100 to 1000 per hour between 2 and 4 AM EDT, so we are dealing with a meteor outburst, potentially a storm. Peak rate of ~200 per hour best current estimate

MAY CAMELOPARDALIDS

On May 24, 2014 Earth will encounter multiple streams of debris laid down by Comet 209P LINEAR

This will produce a new meteor shower, never before seen

Rates predicted to be from 100 to 1000 per hour between 2 and 4 AM EDT, so we are dealing with a meteor outburst, potentially a storm. Peak rate of ~200 per hour best current estimate

Difficult to calibrate models due to lack of past observations

MAY CAMELOPARDALIDS

On May 24, 2014 Earth will encounter multiple streams of debris laid down by Comet 209P LINEAR


This will produce a new meteor shower, never before seen

Rates predicted to be from 100 to 1000 per hour between 2 and 4 AM EDT, so we are dealing with a meteor outburst, potentially a storm. Peak rate of ~200 per hour best current estimate

Difficult to calibrate models due to lack of past observations

Models indicate mm size particles in stream, so potential risk to Earth orbiting spacecraft

Model	Time on May 24 (UT)	ZHR (#/hr)	RA, Dec (°)	V_z (km/s)
<u>Lyytinen & Jenniskens</u> (1929)	3:19			
<u>Lyytinen & Jenniskens</u> (1979)	6:04			
MSFC (peak 1)	6:11			
<u>Lyytinen & Jenniskens</u> (1818, 1853)	6:33			
MSFC (peak 2)	6:56			
<u>Lyytinen & Jenniskens</u> (1903, max)	6:59	15.86	125, +78	15.86
<u>Lyytinen & Jenniskens</u> (1909)	7:15			
<u>Maslov</u> (1898-1919; 1903)	7:18	200-300	122.8, +79.1	16.2
<u>Maslov</u> (max)	7:21	100	122.8, +79.0	
Vaubailon	7:40	100-400	~125, +79	
<u>Lyytinen & Jenniskens</u> (1914)	7:49			
<u>Maslov</u> (1763-1783)	7:55	50-150	122.8, +79.0	16.2
MSFC (peak 3)	8:10			
Jenniskens (general)			123, +79	16





7-year observing program

Goal: Monitor the Moon for impact flashes produced by meteoroids striking the lunar surface.



Observation from MSFC

- Two 0.35m telescopes simultaneously
- Black & white CCD video cameras
- Interleaved 30fps video digitized, recorded
- Video analyzed with custom software




Field of View

- FOV covers approx. 20 arcmin
- 4×10^6 km² on the leading or trailing edge
- Observing when illumination 10-50%
- Maximum 10 observing nights/month

300+ lunar impacts observed

2005-present



For more info:


Suggs et al. (2013) to be submitted

Suggs et al. (2011)


Suggs et al. (2007)

12/15/2006
09:17:39.336

33 ms
 $m_R = 7.4$
0.09 kg
Geminid (35 km/s)



T



Observed by D.E. Moser & W.R. Swift; detected by R.J. Suggs

11/17/2006
10:56:34.820

66 ms
 $m_R = 7.0$
0.03 kg
Leonid (71 km/s)



S




Observed by V. Coffey, D.E. Moser & W.R. Swift; detected by R.J. Suggs

11/03/2008
00:11:06.144

100 ms
 $m_R = 7.7$
0.1 kg
S. Taurid (27 km/s)



T



Observed by W.R. Swift; detected by R.J. Suggs

04/22/2007
03:12:24.372

133 ms
 $m_R = 6.7$
0.08 kg
Lyrid (49 km/s)



T



Observed by R.M. Suggs; detected by R.J. Suggs



Monday, March 31, 14

AVGR - Shot 10

Projectile: 0.25" Pyrex
Target: Pumice Powder
Speed: 5.32 km/s
45 deg. impact angle





AVGR - Shot 10

Projectile: 0.25" Pyrex
Target: Pumice Powder
Speed: 5.32 km/s

011 00:04:712

NASA AVGR

March 17, 2013 3:50:54 UTC

03/17/2013

03:50:54.312

1.03 s


$m_R = ?$

Impactor mass?

Meteor shower association?






Observed by A. Kingery & R.M. Suggs; detected by R.J. Suggs



The brightest meteoroid impact flash detected in NASA's 7 year lunar observing program.




Estimating the magnitude



Saturated

Peak $m_R = 4.9$



2D elliptical Gaussian fit
to the unsaturated wings

Peak $m_R = 3.0 \pm 0.4$

Similar results for
2D elliptical Moffat fit

Photometry
performed using
comparison stars

Preliminary energy estimate




Luminous efficiency (η_λ) relates how much of the impactor's kinetic energy (KE) is converted to luminous energy (LE) in a wavelength range λ

$$LE_\lambda = \eta_\lambda KE_\lambda$$

	Const. $\eta = 2 \times 10^{-4}$		Vel. dep. $\eta = 1.3 \times 10^{-3}$ (Moser et al. 2011)	
	Average	Range	Average	Range
Luminous energy (J)	7.1×10^6	$4.7 \times 10^6 - 1.1 \times 10^7$	7.1×10^6	$4.7 \times 10^6 - 1.1 \times 10^7$
Kinetic energy of impactor (J)	3.6×10^{10}	$2.4 \times 10^{10} - 5.5 \times 10^{10}$	5.4×10^9	$3.6 \times 10^9 - 8.4 \times 10^9$
Impactor mass (kg) (assuming $v_g = 25.6$ km/s)	108	72 – 168	16	11 – 26


Why did we assume $v_g = 25.6$ km/s?





Meteor shower association

19 fireballs seen on Mar 17, 2013




Geocentric meteor radiant color-coded by speed with a tight cluster of 5 with

Virginid Complex			
	meteors	NVI ¹	EVI ²
α_g (°)	184.1 ± 1.0	185.7	183.6
δ_g (°)	4.4 ± 0.9	2.3	3.7
v_g (km/s)	25.6 ± 0.8	23.0	28.9
λ_{sun} (°)	356.6	354	354

¹(Sekanina, 1973), ²(Whipple, 1957)


Cluster of 5 seen on Mar 17, 2013




Orbits of the cluster of 5 were very similar with the following average orbital elements

	meteoroids	NVI	EVI
a (AU)	2.25 ± 0.17	1.69	2.82
e	0.79 ± 0.02	0.71	0.86
i (°)	5.26 ± 1.02	3.7	5.2
ω (°)	280.32 ± 2.11	282.4	285.8
Ω (°)	356.65 ± 0.07	358.0	355.1
q (AU)	0.48 ± 0.02	0.496	0.40
Q (AU)	4.0 ± 0.3	2.89	5.25
Tj	3.1 ± 0.2	→ Indicates asteroidal body	

Mapping the impact location




Flash at peak brightness





Flash 10 frames (333 ms) after the peak

ArcMap (ArcGIS 10) was used to georeference the lunar impact video




Average location: $20.599 \pm 0.172^\circ$ N, $23.922 \pm 0.304^\circ$ W





2014 AA


Discovered January 1 at
6:18 UT. 7 images taken
over 70 minute interval



2014 AA

Discovered January 1 at
6:18 UT. 7 images taken
over 70 minute interval

Magnitude of 19
indicated an object 2-4
meters in diameter





2014 AA

Discovered January 1 at 6:18 UT. 7 images taken over 70 minute interval


Magnitude of 19 indicated an object 2-4 meters in diameter


Positions submitted to the MPC, but no one realized object was on an impact course with Earth until just before impact






Projected impact time was Jan 2 2:25 UTC, just off coast of west Africa





Actual impact time was Jan 2 3:30 UTC +/- 30 minutes



Actual impact time was Jan 2 3:30 UTC +/- 30 minutes


Impact energy of about 1 kiloton

CHELYABINSK

Fireball Properties

Energy	470 kilotons at altitude of 30 km (19 miles) near Chelyabinsk (54.8° N, 61.1° E)
Speed	19 km/s (42,500 mph)
Mass/size	~12,000 tons 19 m (62 ft)
Composition	Ordinary chondrite (LL5)

Comparative Sizes











15.03.2013 17:22:55





Kiloton Airbursts Detected by Infrasound: 2000-2013



A		B	C D E		F G		H I J K L M N O P Q R								S	T	U V		W	X Y Z				
2000			2002			2004	2006								2008		2010		2012					
Energy (kt):	3	1	3	26	1	28	12	1	3	9	1	1	5	6	1	1	2	50	15	5	4	600	2	10